



The AUVI-Q Doctor Discussion Guide

This guide can help you have a productive conversation with your doctor about anaphylaxis and AUVI-Q. Bring it to your child's next appointment to help get the answers you need.

Questions about anaphylaxis

- What are the signs and symptoms of anaphylaxis?
- How can I tell the difference between a mild allergic reaction and a severe allergic reaction?
- What steps should I take if my child experiences a severe allergic reaction?
- Why should I call 911 or seek emergency medical care after a reaction?
- Should I give my child any additional medications if they're having a severe allergic reaction?

Questions about AUVI-Q

- How is AUVI-Q different from an epinephrine pen?
- How many AUVI-Q auto-injectors should we carry with my child?
- Which family members and/or caregivers should know how to use AUVI-Q?
- When is my child old enough to self-carry and self-administer AUVI-Q?
- Why is it important to administer AUVI-Q immediately after an allergic reaction?
- What steps do I need to take after using AUVI-Q?
- What is an anaphylactic emergency plan? Should AUVI-Q be part of that plan?
- Where should I store my child's AUVI-Q?
- Can I get more than one carton of AUVI-Q?

Add your own questions

Indication

AUVI-Q® (epinephrine injection, USP) is a prescription medicine used to treat life-threatening allergic reactions, including anaphylaxis, in people who are at risk for or have a history of serious allergic reactions.

Important Safety Information

AUVI-Q is for immediate self (or caregiver) administration and does not take the place of emergency medical care. Seek immediate medical treatment after using AUVI-Q. Each AUVI-Q contains a single dose of epinephrine. **AUVI-Q should only be injected into your outer thigh, through clothing if necessary.** If you inject a young child or infant with AUVI-Q, hold their leg firmly in place before and during the injection to prevent injuries. Do not inject AUVI-Q into any other part of your body, such as into veins, buttocks, fingers, toes, hands, or feet. If this occurs, seek immediate medical treatment and make sure to inform the healthcare provider of the location of the accidental injection. Only a healthcare provider should give additional doses of epinephrine if more than two doses are necessary for a single allergic emergency. (continued)

Wondering how much you'll pay for AUVI-Q?

Commercial Insurance:

- If your insurance covers AUVI-Q, you'll pay as little as a \$35 copay* for multiple cartons of AUVI-Q when filled in the same transaction
- If your insurance does not cover AUVI-Q, or if you have a high deductible plan, you'll pay no more than \$150. And, if you have a high-deductible plan, you may be eligible to pay \$35 once you've met your prescription deductible

Government Insurance:

- Many government insurance plans cover AUVI-Q with a low copay. Check your coverage

Want to check your AUVI-Q insurance coverage now?

Scan the QR Code or visit www.auvi-q.com/coverage



Talk to your doctor about AUVI-Q

Your doctor can **prescribe AUVI-Q electronically to ASPN Pharmacies** for delivery straight to your doorstep. You can even text "AUVI-Q" to 84790 to have access to the AUVI-Q portal to set up and track your delivery.[†]

[†]Message and data rates may apply.

***Terms and conditions:** Only valid in the United States and DC. Patients with commercial insurance coverage for AUVI-Q may pay a copay of \$35 or less per carton. Maximum savings is up to cost of copay. Actual copay costs may exceed \$35 depending on insurance coverage, plan deductibles, and plan design. OFFER IS NOT INSURANCE. Not valid for prescriptions paid for in part/full by state or federally funded program(s), like Medicare Part D, Medicaid, Vet. Aff., Dept. of Def., or Tricare, and where prohibited by law. Offer cannot be sold, purchased, traded, transferred, or combined with other offers. Cash discount cards are not commercial payers and cannot be used with offer. Kaléo may change offer at any time without notice. Call 844-357-3968 or visit auvi-q.com/terms for offer eligibility questions.

Important Safety Information (continued)

Rarely, patients who use AUVI-Q may develop infections at the injection site within a few days of an injection. Some of these infections can be serious. Call your healthcare provider right away if you have any of the following symptoms at an injection site: redness that does not go away, swelling, tenderness, or the area feels warm to the touch.

If you have certain medical conditions, or take certain medicines, your condition may get worse or you may have more or longer lasting side effects when you use AUVI-Q. Be sure to tell your healthcare provider about all the medicines you take, especially medicines for asthma. Also tell your healthcare provider about all of your medical conditions, especially if you have asthma, a history of depression, thyroid problems, Parkinson's disease, diabetes, heart problems or high blood pressure, have any other medical conditions, are pregnant or plan to become pregnant, or are breastfeeding or plan to breastfeed. Epinephrine should be used with caution if you have heart disease or are taking certain medicines that can cause heart-related (cardiac) symptoms.

Common side effects include fast, irregular or 'pounding' heartbeat, sweating, shakiness, headache, paleness, feelings of over excitement, nervousness, or anxiety, weakness, dizziness, nausea and vomiting, or breathing problems. These side effects usually go away quickly, especially if you rest. Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

Please see the full Prescribing Information and Patient Information at www.auvi-q.com.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call **1-800-FDA-1088**.

The health information contained herein is provided for general educational purposes only. Please consult with your healthcare professional if you have any questions about your health or treatment.

Auvi-Q®
epinephrine injection, USP
0.1 mg/0.15 mg/0.3 mg auto-injectors